



PORTLAND HARBOR RI/FS

**APPENDIX P**

**COMPREHENSIVE BENTHIC APPROACH**

**DRAFT FEASIBILITY STUDY**

**DRAFT**

<p><b>Privileged and Confidential:</b> Work Product Prepared in Anticipation of Litigation</p>
--

March 2015

**Prepared for**  
The Lower Willamette Group

**Prepared by**  
Windward Environmental, LLC

## TABLE OF CONTENTS

TABLE OF CONTENTS .....	I
LIST OF ATTACHMENTS .....	II
LIST OF TABLES .....	III
LIST OF ACRONYMS.....	IV
1.0 COMPREHENSIVE BENTHIC APPROACH .....	1
1.1 Identification of Comprehensive Benthic Risk Areas .....	4
1.2 Results.....	7
2.0 REFERENCES .....	11

## **LIST OF ATTACHMENTS**

- Attachment 1. Mapped Results of the Comprehensive Benthic Approach
- Attachment 2. Data Results of the Comprehensive Benthic Approach

## **LIST OF TABLES**

Table 1.	Contaminants Potentially Contributing to Benthic Risk Based on Predicted Sediment Toxicity LOE
Table 2.	Rationale for Delineation of Each Benthic AOC

## LIST OF ACRONYMS

AOPC	Area of Potential Concern
BERA	Baseline Ecological Risk Assessment
COPC	contaminants of potential concern
DDD	Dichlorodiphenyldichloroethane
DDE	Dichlorodiphenyldichloroethylene
DDT	Dichlorodiphenyltrichloroethane
EPA	U.S. Environmental Protection Agency
FPM	Floating Percentile Model
FS	feasibility study
HCH	Hexachlorocyclohexane
HPAH	High-molecular-weight Polycyclic Aromatic Hydrocarbon
HQ	Hazard Quotient
LOE	Line of Evidence
LPAH	Low-molecular-weight Polycyclic Aromatic Hydrocarbon
LRM	Logistic Regression Model
LWG	Lower Willamette Group
MQ	Mean Quotient
PAH	Polycyclic Aromatic Hydrocarbon
PCB	Polychlorinated Biphenyl
PEC	probable effects concentration
pMax	Maximum probability of toxicity
SMA	sediment management area
SPI	Sediment Profile Imaging
SQG	Sediment Quality Guideline
SQV	sediment quality value
SVOC	Semivolatile Organic Compound
TBT	Tributyltin
total DDx	Sum of all six DDT isomers (2,4'-DDD, 4,4'-DDD, 2,4'-DDE, 4,4' DDE, 2,4'-DDT, and 4,4'-DDT)
TRZ	Toxicity Reference Value
TZW	Transition Zone Water

## **EXECUTIVE SUMMARY**

---

This appendix identifies the areas posing potentially unacceptable risk to the benthic community for use in the draft Feasibility Study (FS) based on the comprehensive benthic risk approach developed by the Lower Willamette Group (LWG) following direction given by the U.S. Environmental Protection Agency (EPA). The appendix details the methods for identifying Comprehensive Benthic Risk Areas and the rationale for the delineation of each of these areas. Maps of the Comprehensive Benthic Risk Areas are included as Attachment 1 to this appendix. The Comprehensive Benthic Risk Areas are part of Sediment Management Area (SMA) and comprehensive remedial alternative development, as explained in Section 5.3 of the draft FS.

## 1.0 COMPREHENSIVE BENTHIC APPROACH

---

The identification of the areas posing potentially unacceptable risk to the benthic community for use in the draft Feasibility Study (FS) was based on the comprehensive benthic approach developed by the Lower Willamette Group (LWG) following direction given by the U.S. Environmental Protection Agency (EPA) in letters dated April 21, 2010 (EPA 2010), and April 4, 2014 (EPA 2014) (see draft FS, Appendix O). In those letters, EPA specified how areas were to be identified and how alternatives were to be evaluated relative to the protection of the benthic community:

- All benthic sediment quality guidelines (SQGs) in the March 24, 2010, list (EPA 2010) will be included in the analysis. If specific SQGs are found to be inconsistent with other lines of evidence (LOEs) listed below, EPA will review the analysis and determine whether these should be included in the draft FS.<sup>1</sup>
- Sediment toxicity bioassays will form the primary LOE for this analysis. The sediment toxicity LOE will include Level 2 (moderate) effects for three endpoints (i.e., *chironomus* [sic] biomass and mortality and *hyalella* [sic] survival) and Level 3 (severe) effects for all four endpoints (i.e., *chironomus* [sic] biomass and mortality and *hyalella* [sic] biomass and mortality) (Shephard 2014).
- The analysis will consider the number and degree of SQG exceedances.
- The analysis will consider other LOEs such as transition zone water (TZW) compared with ambient water quality criteria for the protection of aquatic life and benthic tissue toxicity reference values (TRVs).
- The analysis will consider the presence/absence of nearby sources and examine benthic community structure (e.g., via sediment profile imaging and related information).
- The analysis will consider data quality and data density issues for the SQGs.

In both the Baseline Ecological Risk Assessment (BERA) (Windward 2013) and the draft FS, the primary LOE for identifying benthic community risks was sediment toxicity, represented by survival and growth of the amphipod *Hyalella azteca* and chironomid midge *Chironomus dilutus* in a laboratory setting. When measured toxicity results were not available, toxicity was predicted based on Site-specific sediment quality values (SQVs) derived from multi-variable statistical models (i.e., the Floating Percentile Model [FPM] (Avocet 2003)), Logistic Regression Model [LRM] (Field et al. 1999)], and probable effects concentration [PECs] (MacDonald et al. 2000)).<sup>2</sup> These models estimate the probability of toxicity above a suite of threshold chemical concentrations (i.e., SQVs)

---

<sup>1</sup> The SQVs have subsequently been revised based on additional modeling and negotiations between the LWG and EPA, as documented in Item 11 of Attachment B to a January 12, 2011, LWG letter to EPA (LWG 2011a), the attachment to a February 25, 2011, Remedial Investigation (RI)/FS schedule letter from EPA to the LWG (Humphrey 2011), and the LWG's March 9, 2011, draft response (LWG 2011b) to EPA's February 25, 2011, letter.

<sup>2</sup> See Section 6.2 and Attachment 6 (Part F) of the BERA (Windward 2013) for further information.

derived for the mixture of chemicals found at the Site, or, in the case of PECs, values calculated by third parties from non-Site-specific data and added to the comprehensive benthic approach at EPA's behest.

Because the predictive models are statistical, results are correlative and do not conclusively identify contaminants causing toxicity.<sup>3</sup> All modeling approaches were used to identify contaminants whose sediment concentrations, when considered in aggregate, appear to help explain the observed toxicity and to identify threshold concentrations for each contaminant above which toxicity was likely to occur (Table 1).<sup>4</sup>

**Table 1. Contaminants Potentially Contributing to Benthic Risk Based on Predicted Sediment Toxicity LOE**

Contaminant	
<b>Metals</b>	
Cadmium	Lead
Chromium <sup>a</sup>	Mercury <sup>a</sup>
Copper	Silver
<b>PAHs</b>	
2-Methylnaphthalene	Dibenzo(a,h)anthracene
Acenaphthene	Fluoranthene
Acenaphthylene	Fluorene
Anthracene	Indeno(1,2,3-cd)pyrene
Benzo(a)anthracene	Phenanthrene
Benzo(b)fluoranthene	Pyrene
Benzo(b+k)fluoranthene	Total HPAHs
Benzo(g,h,i)perylene	Total LPAHs <sup>a</sup>
Benzo(k)fluoranthene	Total PAHs
Chrysene	
<b>Phthalates</b>	
Dibutyl phthalate	
<b>SVOCs</b>	

<sup>3</sup> Risk conclusions based on the secondary benthic LOEs—tissue residue, surface water, and TZW—identify contaminants posing potentially unacceptable risk (i.e., contaminants of potential concern [COPCs]) and are noted in Sections 12.1 and 12.2 of the draft final BERA (Windward 2013).

<sup>4</sup> The contaminant list is a combination of SQVs derived using the FPM and the LRM. Each SQV has a different reporting basis depending on the normalization selected for the model. All FPM SQVs were dry-weight normalized. LRM SQVs used a number of different normalizations, including dry weight, organic carbon, percent fines, and combinations of normalizations.



**Table 1. Contaminants Potentially Contributing to Benthic Risk Based on Predicted Sediment Toxicity LOE**

Contaminant	
Benzyl alcohol	Dibenzofuran <sup>a</sup>
1,2-Dichlorobenzene	Carbazole <sup>a</sup>
<b>Phenols</b>	
4-Methylphenol <sup>b</sup>	Phenol
<b>PCBs</b>	
Total PCBs <sup>a</sup>	
<b>Pesticides</b>	
2,4'-DDD	beta-HCH
4,4'-DDD	delta-HCH <sup>a</sup>
4,4'-DDE	Dieldrin
4,4'-DDT	Endrin
Sum DDD <sup>a</sup>	Endrin ketone
Sum DDE	cis-Chlordane
Sum DDT	Total endosulfan <sup>b</sup>
Total DDx	
<b>Petroleum Hydrocarbons</b>	
Diesel-range hydrocarbons	

**Notes:**

- <sup>a</sup> FPM SQVs based on one or two endpoints are less than the apparent effect threshold and therefore might contribute to false predictions of toxicity.
- <sup>b</sup> All SQVs derived from the FPM are less than the apparent effect threshold and therefore may contribute to false predictions of toxicity.

DDD – dichlorodiphenyldichloroethane  
DDE – dichlorodiphenyldichloroethylene  
DDT – dichlorodiphenyltrichloroethane  
FPM – floating percentile model  
HCH – hexachlorocyclohexane  
HPAH – high-molecular-weight polycyclic aromatic hydrocarbon

LOE – line of evidence  
LPAH – low-molecular-weight polycyclic aromatic hydrocarbon  
PAH – polycyclic aromatic hydrocarbon  
PCB – polychlorinated biphenyl  
SQV – sediment quality value  
SVOC – semivolatile organic compound  
total DDx – sum of all six DDT isomers (2,4'-DDD, 4,4'-DDD, 2,4'-DDE, 4,4'-DDE, 2,4'-DDT, and 4,4'-DDT)

## 1.1 IDENTIFICATION OF COMPREHENSIVE BENTHIC RISK AREAS

Because the primary benthic LOE (bioassay results) does not identify the cause of the empirical toxicity, delineating areas posing potentially unacceptable risk to the benthic

community based on the magnitude of single chemical concentrations is highly uncertain. Rather, the draft FS focused on the empirical evidence of toxicity, along with predictions of toxicity (exceedances of a suite of SQVs derived from three models), to identify spatial aggregations representing areas that pose potentially unacceptable risk to the benthic community (i.e., comprehensive benthic risk areas). A weight of evidence framework that combined the frequency and magnitude of empirical toxicity, frequency and magnitude of toxicity predictions, concordance between models and endpoints, results from other LOEs (i.e., benthic tissue burdens and TZW water quality exceedances), and spatial distribution of endpoints indicating potentially unacceptable risk composed the comprehensive benthic approach. Because surface water quality is not location specific, it was not used to delineate comprehensive benthic risk areas (although it was used to confirm the contribution of specific chemicals to areas posing potentially unacceptable risk to the benthic community in specific reaches of the Site). Sediment profile imaging (SPI) data were not used in the development of comprehensive benthic risk areas because the information was qualitative and the SPI results did not identify any additional comprehensive benthic risk areas; rather, the SPI results generally indicated that benthic community structure could be explained by physical habitat characteristics and hydrological regime (Windward 2013).

Comprehensive benthic risk areas were identified based on the application of the comprehensive benthic approach. Maps are included herein as Attachment 1 to this appendix.

Locations where empirical bioassay results indicated significant toxicity formed the core of a comprehensive benthic risk area. Predictions of toxicity or bioaccumulation at surrounding chemistry-only stations were used as part of the weight of evidence that the area posed potentially unacceptable risk to the benthic community. In areas where no empirical bioassay data were available, predicted toxicity was sufficient to identify comprehensive benthic risk areas. Significant bioaccumulation in either field-collected or laboratory-exposed organisms provided an independent LOE to either corroborate the identification of a comprehensive benthic risk area or provide other evidence that a comprehensive benthic risk area was present.

TZW locations with chemical concentrations that, because of magnitude, were unlikely to be addressed by source control alone, also contributed to the identification of comprehensive benthic risk areas.

Sediment chemistry used to predict toxicity was represented as an aggregate value based on average exceedance factors (i.e., mean quotients [MQs]) across the entire FPM and PEC SQV sets. This MQ method of quantifying the concentrations of multiple chemicals that may be contributing to the potentially unacceptable risk to the benthic community has been used at other sites throughout the United States and was required by EPA in the problem formulation in the BERA (Windward 2013). Predictions of toxicity based on the LRM were represented as the maximum probability of toxicity (pMax) across all chemicals with some potential contribution to the toxicity observed in the empirical

bioassays. Decision thresholds selected by EPA for use in the BERA (Windward 2013) were retained for the comprehensive benthic approach (i.e., an MQ greater than or equal to 0.7 for both LRM and PEC models and a pMax value greater than or equal to 0.59 for the LRM model).

Details of the approach used to identify comprehensive benthic risk areas are as follows:

1. Areas of potential concern (AOPCs) based on multiple LOEs (i.e., benthic community, fish, wildlife, and human health endpoints) were developed by EPA prior to submittal of the BERA (Windward 2013).
2. Locations within these broader AOPCs with empirical bioassay results indicating significant toxicity were identified.
  - a. Significant toxicity was considered to be one toxicity endpoint exceeding a Level 3 threshold, or two endpoints exceeding a Level 2 threshold (Shephard 2014).
    - i. Level 3 threshold: Four empirical toxicity test endpoints (i.e., *Chironomus* biomass or growth or *Hyaella* biomass or growth).
    - ii. Level 2 threshold: Three empirical toxicity test endpoints (i.e., *Chironomus* biomass or growth or *Hyaella* survival)
3. Locations without bioassay data but where significant sediment toxicity was predicted based on sediment chemistry exceedances were identified.
  - a. Sampling locations where at least two of the three models' thresholds were exceeded were considered toxic.
  - b. Sampling locations where no threshold or only one threshold was exceeded were considered non-toxic.
4. Locations where empirical tissue residues or predicted tissue residues (when empirical tissue residue data were absent) exceeded their TRVs were identified.
  - a. The evidence of risk provided by measured or predicted exceedance of metals TRVs was considered weak because of species-specific differences in metals sequestration or other bioregulation; such evidence was not used to identify comprehensive benthic risk areas.
  - b. The evidence of risk provided by a predicted exceedance of the tributyltin (TBT) TRV was considered weak because of high uncertainty in the TBT bioaccumulation model and the selected TRV. Bioaccumulation (predicted or measured) of this chemical was not used to identify comprehensive benthic risk areas.
5. TZW exceedance areas with hazard quotients (HQs) greater than 10 were delineated separately (see Section 6.6.3.3 of the BERA (Windward 2013) for explanation of why 10 is a conservative threshold for the TZW HQ).
6. Individual sample results representing each benthic LOE were overlaid on a map.

- a. Comprehensive benthic risk areas were identified where two or more adjacent sampling locations indicated potentially unacceptable risk to the benthic community based on either empirical or predicted toxicity, empirical or predicted bioaccumulation, empirical TZW chemistry, or a combination of bioassay and chemistry LOEs.
  - i. Because empirical toxicity was the primary LOE, toxicity predicted by chemistry exceedances (i.e., MQs or pMax) were overridden by no-hit bioassays where these lines co-occurred.
- b. TZW exceedance areas (based on HQs greater than 10) were identified as comprehensive benthic risk areas.
- c. Boundaries of the comprehensive benthic risk areas split the distance between sampling locations exceeding criteria and the surrounding clean sampling locations, except where:
  - i. Other physical features (e.g., pier, channel edge, property boundary) were present, in which case the boundaries were drawn at the physical features.
  - ii. The nearest sampling location exceeding criteria was at a distance greater than 200 feet<sup>5</sup> from a clean sampling location, in which case the boundary was drawn at a subjective distance less than halfway to the nearest clean sampling location.

## 1.2 RESULTS

The results of the application of the comprehensive benthic approach are presented in Attachments 1 and 2 to this appendix. Table 2 summarizes the rationale for the delineation of a comprehensive benthic risk area.

**Table 2. Rationale for Delineation of Each Comprehensive Benthic Risk Area**

AOC	Is it a Comprehensive Benthic Risk Area?	Rationale
1A-1	Yes	Five locations with empirical and predicted bioaccumulation above TRVs were clustered together along the shoreline.
1A-2	No	Only two bioassay and one chemistry location exceeded their respective thresholds. Two predicted bioaccumulation locations were not contiguous with the risk area.
2	No	Three bioassay Level 2 or 3 hits. No other LOE indicating toxicity.

<sup>5</sup> This distance is based on best professional judgment. Conditions within Portland Harbor tend to be localized with few apparent gradients. In addition, sampling density varied with mid-channel samples being the least dense, necessitating selection of some distance to delineate a cluster of samples that had some likelihood of being associated with similar sources.

**Table 2. Rationale for Delineation of Each Comprehensive Benthic Risk Area**

<b>AOC</b>	<b>Is it a Comprehensive Benthic Risk Area?</b>	<b>Rationale</b>
3-1	No	Seven locations with significant toxicity or actual or predicted bioaccumulation were spatially isolated.
3-2	Yes	Twelve locations with significant toxicity and bioaccumulation (either empirical or predicted) formed a cluster.
4	No	Single, isolated bioassay Level 3 hit with no other LOE indicating toxicity.
5-1	Yes	Three empirical toxicity locations formed a cluster.
5-2	No	Two bioassay locations indicating toxicity were spatially isolated.
6-1	Yes	Eight locations with predicted toxicity formed a cluster.
6-2	No	Two locations with empirical or predicted toxicity; locations were spatially isolated.
7	No	No LOE indicating toxicity.
8-1	No	Two locations with significant toxicity were spatially isolated.
9D-1	No	Fourteen locations with significant empirical or predicted toxicity or predicted bioaccumulation were spatially separated by non-toxic locations.
9D-2	Yes	Three locations with empirical or predicted toxicity formed a cluster.
9D-3	Yes	Five locations with predicted toxicity formed a cluster.
9U-1a	Yes	Twenty-six locations with significant toxicity, bioaccumulation, and TZW HQs greater than 100 formed a cluster.
9U-1b	Yes	Fifty-one locations with significant toxicity, bioaccumulation, and TZW HQs greater than 10 formed a cluster.
9U-2	No	No LOE indicating toxicity.
10	No	Two stations indicating empirical toxicity; locations were spatially isolated.
11	No	Single isolated prediction of toxicity; no other LOE indicating toxicity.
12	No	Single isolated prediction of toxicity; no other LOE indicating toxicity.
13-1	No	Eight locations with significant empirical or predicted toxicity or empirical bioaccumulation were spatially separated by non-toxic locations.
13-2	Yes	Five locations with empirical or predicted toxicity or bioaccumulation formed a cluster.
13-3	Yes	Two locations with empirical or predicted toxicity were adjacent to a previously remediated area.
14-1	No	Six locations with significant empirical or predicted toxicity or bioaccumulation were spatially separated by non-toxic locations.

**Table 2. Rationale for Delineation of Each Comprehensive Benthic Risk Area**

<b>AOC</b>	<b>Is it a Comprehensive Benthic Risk Area?</b>	<b>Rationale</b>
14-2	Yes	Two locations with empirical or predicted toxicity formed a small cluster.
14-3	Yes	Fourth-three locations with significant toxicity, bioaccumulation, and TZW HQs greater than 10 formed a cluster.
14-4	Yes	Two locations with empirical or predicted toxicity formed a small cluster.
15	No	No LOE indicating toxicity.
16	No	Four locations with empirical and predicted toxicity were spatially separated by intervening clean locations.
17D-1	No	Nine locations with either toxicity or predicted bioaccumulation were spatially isolated.
17S-1	No	Locations with significant toxicity or bioaccumulation were spatially separated.
17S-2	Yes	Ten locations with either empirical or predicted toxicity or bioaccumulation formed a cluster.
17S-3	Yes	Three locations with predicted toxicity or bioaccumulation formed a cluster.
18	No	Six locations with predicted and empirical toxicity were spatially isolated.
19-1	No	Four locations with significant predicted and empirical toxicity were spatially isolated.
19-2	Yes	Eleven locations with empirical and predicted toxicity and bioaccumulation locations formed a cluster.
19-3	Yes	Five locations with empirical and predicted toxicity and predicted bioaccumulation locations formed a cluster.
20	No	Two locations, one isolated bioaccumulation location and the other one empirical toxicity location were spatially isolated.
21	No	No LOE indicating toxicity.
22	No	No LOE indicating toxicity.
23	No	One location with empirical toxicity was spatially isolated.
24	No	No LOE indicating toxicity.
25-1	No	Two locations indicating significant bioaccumulation were spatially isolated.
25-2	Yes	Eight locations with predicted toxicity and six bioaccumulation locations formed a cluster.
26	No	No LOE indicating toxicity.

AOC – area of concern

TRV – toxicity reference value

HQ – hazard quotient  
LOE – line of evidence

TZW – transition zone water

Eighteen comprehensive benthic risk areas were identified based on sediment toxicity, bioaccumulation, and TZW LOEs. One comprehensive benthic risk area (1A-1) was identified based solely on the bioaccumulation LOE. Six comprehensive benthic risk areas (9D-2, 9U-1a, 13-3, 14-2, 14-4, 17s-2) were identified based on toxicity, both predicted and empirical. One comprehensive benthic risk area (5-1) was identified based solely on empirical bioassay results, while two others (6-1 and 9D-3) were identified based solely on predicted toxicity. Three comprehensive benthic risk areas (9U-1a, 9U-1b, and 14-3) had TZW TRV HQs exceeding 10; however, other LOEs also indicated toxicity in these areas.

## 2.0 REFERENCES

---

- Avocet. 2003. Development of freshwater sediment quality values for use in Washington State. Phase II report: Development and recommendation of SQVs for freshwater sediments in Washington State. Publication No. 03-09-088. Prepared for Washington Department of Ecology. Avocet Consulting, Kenmore, WA.
- EPA. 2010. EPA letter and attachment dated April 21, 2010 to Lower Willamette Group (from E. Blischke and C. Humphrey to J. McKenna) regarding Portland Harbor Superfund site: EPA direction to LWG on preliminary remediation goals (PRGs) for use in the Portland Harbor feasibility study. US Environmental Protection Agency Region 10, Oregon Operations Office, Portland, OR.
- Field LJ, MacDonald DD, Norton SB, Severn CG, Ingersoll CG. 1999. Evaluating sediment chemistry and toxicity data using logistic regression modeling. *Environ Toxicol Chem* 18:1311-1322.
- Humphrey C. 2011. Personal communication (re: Portland Harbor AOC for RI/FS Schedule). US Environmental Protection Agency, Seattle, WA. February 25, 2011.
- LWG. 2011a. Letter and attachments dated January 12, 2011 from B. Wyatt to C. Humphrey, EPA Region 10: response to December 21, 2010 EPA letter on the status of the Portland Harbor Feasibility Study; September 27, 2010 EPA letter on the Benthic Risk Evaluation; and December 8, 2010 EPA letter on general responses to EPA non-directed RI, BHHRA and BERA comments. Lower Willamette Group, Portland, OR.
- LWG. 2011b. Letter dated March 9, 2011, from Bob Wyatt to Chip Humphrey and Kristine Koch, EPA Region 10: response to EPA's February 25, 2011, response to LWG proposed project schedule presented on February 2, 2011. Draft. Lower Willamette Group, Portland, OR.
- MacDonald DD, Ingersoll CG, Berger TA. 2000. Development and evaluation of consensus-based sediment quality guidelines for freshwater ecosystems. *Arch Environ Contam Toxicol* 39(5):20-31.
- Shephard B. 2014. Personal communication (email from B. Shephard, EPA, to J. Toll, Windward, regarding comprehensive benthic approach thoughts and recommendations for the FS). US Environmental Protection Agency, Seattle, WA. April 4, 2014.
- Windward. 2013. Portland Harbor RI/FS, Final Remedial Investigation Report, Appendix G: Baseline ecological risk assessment. Final. Prepared for the Lower Willamette Group. Windward Environmental LLC, Seattle, WA.



# **Attachment 1**

## **Mapped Results of the Comprehensive Benthic Approach**

## **Attachment 2**

# **Data Results of the Comprehensive Benthic Approach**